3<sup>rd</sup> International Conference on

## NURSING & MIDWIFERY

May 23-24, 2018 | New York, USA

## Quality of life of patients undergoing cancer treatment in B.P. Koirala Memorial Cancer Hospital, Bharatpur, Chitwan

Radha Acharya Pandey<sup>1</sup>, Dhungana Govinda Prasad<sup>2</sup>, Jyotsana Twi Twi<sup>1</sup>, Sudeepa Byanju<sup>1</sup> and Barsha Khawas<sup>1</sup> <sup>1</sup>Kathmandu University School of Medical Sciences, Nepal <sup>2</sup>Chitwan Medical College, Nepal

**Abstract:** This research entitled "Quality of Life of Patients Undergoing Cancer Treatment in B.P. Koirala Memorial Cancer Hospital, Bharatpur, Chitwan. It was conducted to assess the quality of life of cancer patients. It was carried out among patients attending B. P. Koirala Memorial Cancer Hospital, Bharatpur, Chitwan.

**Background:** In patients with different type of cancers and the quality of life (QoL) improvement is the main goal, since survival can be prolonged marginally. A diagnosis is very stressful for people, affecting all aspects of their being and quality of life. Up to date, knowledge on QoL impairments throughout the entire treatment process, often including several treatment modalities is scarce. One objective of this study was to assess the quality of life of cancer patient undergoing cancer treatment.

**Methods:** A quantitative, cross-sectional, descriptive, design was adapted. The total of 200 and 45 cancer patients met the inclusion criteria and were enrolled in the studies who were attending the hospital for cancer treatment during August-September, 2013. The data was collected by interview, using modified, structured scale of European Organization for Research and Treatment of Cancer Quality of life Questionnaire (EORTC QLQ- C30), prepared by the EORTC group. Information about the patient's disease condition and treatment were obtained from the patient's medical records. The collected data was analyzed by using SPSS version 16. Descriptive and inferential statistics were used to describe the respondent's quality of life (QoL) scores and to identify the factors affecting it respectively.

**Results:** The study findings revealed the quality of life of cancer patients to be influenced by many factors such as: site of cancer, stage of cancer, time elapsed since diagnosis and Eastern Co-operative Oncology Group (ECOG) performance status. The average QoL scores (out of 100) for different scales were 85.54 (global health/QoL), 77.03 (functional), and 16.14 (symptom). Loss of appetite was the most frequent complaint (mean=20.27) and was present in almost all the patients. As the overall, QoL of the patients was significantly correlated with different QoL scales as, cognitive, emotional, physical, social, role functioning, pain, fatigue, dyspnoea, loss of appetite and nausea/vomiting and financial problem.

**Conclusion:** Hence, in average, the quality of life of cancer patients was found to be relatively better, although there were higher ratings for some (as: cognitive, physical, role and emotional functioning) and lower for others (like social functioning). Additional research should be done in this area for improving the quality of life of specific type of cancer patients in Nepal, though the findings of this study are expected to provide the baseline knowledge regarding it.

## Biography

Radha Acharya Pandey has completed her Master degree in Adult Nursing from Institute of Medicine, Nepal. She is an Assistant Professor of Kathmandu University School of Medical Sciences. She published more than 15 papers in reputed journals and she served as a Reviewer for a number of journals in her related field and earned certificate for quality review. She has presented paper at international and national conferences and organized conference, workshop in related areas.

radhapnd@yahoo.com

Notes: