

3rd International Conference on

NURSING & MIDWIFERY

May 23-24, 2018 | New York, USA

The evidence-based C-section its overuse and how midwives can contribute to decrease its rate

Michael Stark

New European Surgical Academy, Germany

As most abdominal operations have endoscopic alternatives, caesarean section will remain the only abdominal operation in the future. Therefore it is of utmost importance to constantly evaluate the different steps for their necessity and for their optimal way of performance. The modified Joel-Cohen method results in a shorter incision to delivery time, lower rate of febrile morbidity compared to the traditional Pfannenstiel incision. Opening peritoneum using bi-digital stretching rather than sharp instruments proved to be safer, and exteriorization of the uterus makes stitching easier and avoids unnecessary bleeding. Suturing the uterus with one layer only results in stronger scars and reduced pain. Leaving both peritoneum layers open reduces adhesions. The fascia being sutured continuously with first knot underneath the fascia prevents irritation in the sub-cutis, and, by a right-handed surgeon, from the right to the left, proved to be ergonomic. Since the introduction of this modified and simplified method, it has been evaluated in dozens of peer-reviewed publications from different countries. Without exception, all showed various advantages of this method: shorter operation time, shorter hospitalization, quicker mobilization, less blood loss, lower rate of febrile morbidity, lower costs, and less need for painkillers. Only 10 instruments and three sutures are needed, which simplifies the workload of nurses. In order to standardize this operation, it is important to use constantly the same needles and instruments. Big needle is necessary for the uterus, as fewer steps are done and therefore less foreign body reaction. This operation is recommended as universal routine method for caesarean section and its principles should apply to all surgical disciplines. Unfortunately, the rate of cesarean section is rising constantly around the world. As evolution continues, it might be influenced by this high rate. In this presentation, the logic of the need to limit the numbers of cesarean section based on anthropological studies will be presented as well as the role of midwives in decreasing its rate through observation, communication between parturient and obstetrician and attention to physiological processes. NESAs anti-caesarean project will be introduced.

mstark@nesacademy.org

Notes: