

## Risk factors for fast early infarct growth in patients with acute ischemic stroke

**Han Liu**

JiulongPo District People's Hospital, Chongqing, China

**Background:** Fast early infarct growth (EIG) has been shown to be an important predictor of patient neurological outcomes and all-cause mortality. However, the factors related to fast EIG remain largely unknown. The present study aimed to elucidate the risk factors for fast EIG in acute ischemic stroke with a large vessel occlusion.

**Materials and Methods:** In this single-center prospective study, patients with acute ischemic stroke due to anterior circulation large vessel occlusion identified by CT or MRI 24-h from onset were included and analyzed. Fast EIG was defined as infarct volume on baseline imaging/onset-to-imaging time  $\geq 10$  mL/h..

**Results:** Overall, 149 patients, including 83 males and 56 females at a median age of 71 years and median NIS Stroke Scale of 15 (interquartile range 8-22) were included. A total of 58 (38.93%) patients were determined to have fast EIG. A higher age (HR=2.42, P=0.009), advanced NIS Stroke Scale (HR=1.43, P=0.02) and poor collateral perfusion on imaging (HR=5.21, P<0.001) were noted to be independent risk factors for a fast EIG. In addition, patients with fast EIG had significantly unfavorable neurological outcomes as indicated by the 3-month modified Rankin Scale than those without fast EIG (P<0.001).

**Conclusions:** Advanced age, NIS Stroke scale and poor collateral perfusion were risk factors for fast EIG in acute ischemic patients with a large vessel occlusion.

### Biography

Han Liu, JiulongPo District People's Hospital, Chongqing, China

**Received:** September 17, 2024; **Accepted:** September 18, 2024; **Published:** December 13, 2024