

Investigating the Molecular Interactions between FGFR2 and NOTCH1 Pathways: Implications for Targeted Cancer Therapies and Treatment Strategies

Santiago Antonio*

Department of Pathology, School of Medicine, National and Kapodistrian University of Athens, 11527 Athens, Greece

Introduction

Among the many signaling pathways implicated in tumorigenesis, the Fibroblast Growth Factor Receptor 2 (FGFR2) and NOTCH1 pathways are two key contributors to the development and progression of several cancers. Recent research has highlighted the interactions between these two pathways, revealing their combined role in regulating processes like cell proliferation, survival, migration, and metastasis. Understanding these interactions has profound implications for the development of targeted cancer therapies that can specifically target these pathways to treat various malignancies, including breast, lung, and gastric cancers. Cancer remains one of the most formidable health challenges globally, with molecular interactions within cells often playing critical roles in its development and progression. Among these interactions, those involving Fibroblast Growth Factor Receptor 2 (FGFR2) and NOTCH1 have garnered significant attention. Both FGFR2 and NOTCH1 are crucial signalling molecules that regulate various cellular processes, including proliferation, differentiation and survival. This article delves into the molecular interactions between FGFR2 and NOTCH1, exploring their implications for cancer therapy.

Description

FGFR2 is a member of the fibroblast growth factor receptor family, which consists of receptor tyrosine kinases. These receptors are activated by binding to fibroblast growth factors (FGFs), leading to receptor dimerization and auto phosphorylation. This activation triggers downstream signalling pathways, such as the MAPK/ERK, PI3K/AKT and PLC pathways, which are involved in cell proliferation, survival and differentiation. Aberrant FGFR2 signaling has been implicated in various cancers, including breast, gastric and lung cancers. Mutations, amplifications and fusions of the FGFR2 gene can lead to its constitutive activation, promoting oncogenesis. Targeting FGFR2 with specific inhibitors has shown promise in clinical trials, highlighting its potential as a therapeutic target. The NOTCH signaling pathway is a highly conserved cell signalling system present in most multicellular organisms. NOTCH1, one of the four NOTCH receptors in mammals, is activated by binding to its ligands (Delta-like and Jagged families). Upon ligand binding, NOTCH1 undergoes photolytic cleavage, releasing the intracellular domain (NICD). The NICD translocates to the nucleus, where it regulates the transcription of target genes involved in cell fate determination, differentiation and proliferation [1].

NOTCH1 plays a dual role in cancer, acting as either an oncogene or a tumour suppressor, depending on the context. In certain cancers, such as T-cell acute lymphoblastic leukaemia (T-ALL), NOTCH1 is frequently mutated, leading to its activation and promoting cancer cell survival and proliferation.

***Address for Correspondence:** Santiago Antonio, Department of Pathology, School of Medicine, National and Kapodistrian University of Athens, 11527 Athens, Greece; E-mail: antonio@santiago.uoa.gr

Copyright: © 2024 Antonio S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 26 October, 2024, Manuscript No. jjo-24-156567; **Editor Assigned:** 28 October, 2024, PreQC No. P-156567; **Reviewed:** 11 November, 2024, QC No. Q-156567; **Revised:** 18 November, 2024, Manuscript No. R-156567; **Published:** 25 November, 2024, DOI: 10.37421/2329-6771.2024.13.518

Conversely, in other cancers, NOTCH1 functions as a tumour suppressor and its loss contributes to tumour genesis. Emerging evidence suggests a complex interplay between FGFR2 and NOTCH1 signalling pathways. This crosstalk can influence various cellular outcomes and has significant implications for cancer therapy. FGFR2 activation can influence the expression of NOTCH1 ligands, thereby modulating NOTCH1 signalling. Conversely, NOTCH1 signalling can affect the expression of FGFR2 and its downstream targets. Both FGFR2 and NOTCH1 pathways converge on common downstream effectors, such as the MAPK/ERK pathway. Crosstalk at this level can amplify or dampen signalling outputs, affecting cell fate decisions. FGFR2 and NOTCH1 can undergo post-translational modifications, such as phosphorylation and ubiquitination that affect their stability and activity. Crosstalk between these pathways can modulate these modifications, altering the signalling dynamics [2,3].

Understanding the molecular interactions between FGFR2 and NOTCH1 provides new avenues for therapeutic intervention. Several strategies can be envisioned. Targeting both FGFR2 and NOTCH1 pathways simultaneously may produce synergistic effects, enhancing therapeutic efficacy and overcoming resistance to single-agent therapies. Identifying biomarkers that reflect the status of FGFR2 and NOTCH1 signalling can guide the selection of patients who are most likely to benefit from targeted therapies. Given the context-dependent roles of FGFR2 and NOTCH1 in cancer, therapies need to be tailored to the specific genetic and molecular landscape of each tumour. Several FGFR2 inhibitors are currently in clinical development or have received regulatory approval. These include small molecule tyrosine kinase inhibitors, monoclonal antibodies and FGF ligand traps. Clinical trials have shown promising results, particularly in cancers with FGFR2 alterations. NOTCH1 inhibitors, including -secretase inhibitors (GSIs) and monoclonal antibodies targeting NOTCH1 ligands, have been investigated in preclinical and clinical studies. While these inhibitors have shown efficacy in certain cancers, challenges such as toxicity and resistance need to be addressed.

Despite the promising therapeutic potential of targeting FGFR2 and NOTCH1, several challenges remain. Cancer cells can develop resistance to FGFR2 and NOTCH1 inhibitors through various mechanisms, such as secondary mutations and activation of compensatory pathways. Understanding these resistance mechanisms is crucial for developing effective combination therapies. Inhibition of FGFR2 and NOTCH1 signalling can lead to adverse effects, given their roles in normal tissue homeostasis. Strategies to minimize toxicity while maintaining therapeutic efficacy are needed. Reliable biomarkers are essential for patient stratification and monitoring therapeutic responses. Advances in genomics and proteomics hold promise for identifying such biomarkers [4,5].

Conclusion

The molecular interplay between FGFR2 and NOTCH1 pathways plays a critical role in the regulation of tumor growth, metastasis, and resistance to therapy. Their interactions contribute to key processes like cancer stem cell maintenance, angiogenesis, and epithelial-to-mesenchymal transition, all of which are critical for tumor progression. Understanding these pathways' cross-talk provides new opportunities for developing targeted therapies aimed at disrupting these mechanisms. Dual inhibition strategies, cancer stem cell targeting, and combination therapies hold great promise for improving outcomes in cancers driven by aberrant FGFR2 and NOTCH1 signaling. As research progresses, these findings could lead to more effective, personalized

treatment strategies for patients with cancers dependent on these pathways. The intricate molecular interactions between FGFR2 and NOTCH1 have profound implications for cancer therapy. By elucidating the mechanisms of crosstalk and developing targeted interventions, we can improve therapeutic outcomes for patients with cancers driven by aberrant FGFR2 and NOTCH1 signalling. Future research should focus on overcoming the challenges of resistance, toxicity and biomarker identification to fully realize the potential of targeting these pathways in cancer therapy.

Acknowledgement

None.

Conflict of Interest

None.

References

1. Tripathy, Nirmalya, Rafiq Ahmad, Seung Hyuck Bang and Gilson Khang, et al. "Outstanding antibiofilm features of quanta-CuO film on glass surface." *ACS Appl Mater Interfaces* 8 (2016): 15128-15137.
2. Xiang, Fan, Robyn Lucas, Simon Hales and Rachel Neale, et al. "Incidence of nonmelanoma skin cancer in relation to ambient UV radiation in white populations, 1978-2012: Empirical relationships." *JAMA Dermatol* 150 (2014): 1063-1071.
3. Kaufman, Howard L., Kim Margolin and Ryan Sullivan. "Management of metastatic melanoma in 2018." *JAMA Oncol* 4 (2018): 857-858.
4. Volpe, Virginia O., Daniel M. Klufas, Upendra Hegde and Jane M. Grant-Kels, et al. "The new paradigm of systemic therapies for metastatic melanoma." *J Am Acad Dermatol* 77 (2017): 356-368.
5. Ugurel, Selma, Joachim Rohmel, Paolo A. Ascierto and Keith T. Flaherty, et al. "Survival of patients with advanced metastatic melanoma: The impact of novel therapies—update 2017." *Eur J Cancer* 83 (2017): 247-257.

How to cite this article: Antonio, Santiago. "Investigating the Molecular Interactions between FGFR2 and NOTCH1 Pathways: Implications for Targeted Cancer Therapies and Treatment Strategies." *J Integr Oncol* 13 (2024): 518.