

Molecular Interactions between FGFR2 and NOTCH1: Implications for Cancer Therapy

Santiago Antonio*

Department of Pathology, School of Medicine, National and Kapodistrian University of Athens, 11527 Athens, Greece

Introduction

Cancer remains one of the most formidable health challenges globally, with molecular interactions within cells often playing critical roles in its development and progression. Among these interactions, those involving Fibroblast Growth Factor Receptor 2 (FGFR2) and NOTCH1 have garnered significant attention. Both FGFR2 and NOTCH1 are crucial signaling molecules that regulate various cellular processes, including proliferation, differentiation and survival. This article delves into the molecular interactions between FGFR2 and NOTCH1, exploring their implications for cancer therapy.

Description

FGFR2 signaling pathway

FGFR2 is a member of the fibroblast growth factor receptor family, which consists of receptor tyrosine kinases. These receptors are activated by binding to fibroblast growth factors (FGFs), leading to receptor dimerization and autophosphorylation. This activation triggers downstream signaling pathways, such as the MAPK/ERK, PI3K/AKT and PLC γ pathways, which are involved in cell proliferation, survival and differentiation [1,2].

FGFR2 in cancer

Aberrant FGFR2 signaling has been implicated in various cancers, including breast, gastric and lung cancers. Mutations, amplifications and fusions of the FGFR2 gene can lead to its constitutive activation, promoting oncogenesis. Targeting FGFR2 with specific inhibitors has shown promise in clinical trials, highlighting its potential as a therapeutic target [3].

NOTCH1 signaling pathway

The NOTCH signaling pathway is a highly conserved cell signaling system present in most multicellular organisms. NOTCH1, one of the four NOTCH receptors in mammals, is activated by binding to its ligands (Delta-like and Jagged families). Upon ligand binding, NOTCH1 undergoes proteolytic cleavage, releasing the intracellular domain (NICD). The NICD translocates to the nucleus, where it regulates the transcription of target genes involved in cell fate determination, differentiation and proliferation.

NOTCH1 in cancer

NOTCH1 plays a dual role in cancer, acting as either an oncogene or a tumor suppressor, depending on the context. In certain cancers, such as T-cell acute lymphoblastic leukemia (T-ALL), NOTCH1 is frequently mutated,

leading to its activation and promoting cancer cell survival and proliferation. Conversely, in other cancers, NOTCH1 functions as a tumor suppressor and its loss contributes to tumorigenesis [4].

Molecular crosstalk between FGFR2 and NOTCH1

Emerging evidence suggests a complex interplay between FGFR2 and NOTCH1 signaling pathways. This crosstalk can influence various cellular outcomes and has significant implications for cancer therapy.

Mechanisms of Interaction

- Regulation of gene expression:** FGFR2 activation can influence the expression of NOTCH1 ligands, thereby modulating NOTCH1 signaling. Conversely, NOTCH1 signaling can affect the expression of FGFR2 and its downstream targets.
- Shared downstream effectors:** Both FGFR2 and NOTCH1 pathways converge on common downstream effectors, such as the MAPK/ERK pathway. Crosstalk at this level can amplify or dampen signaling outputs, affecting cell fate decisions.
- Post-translational modifications:** FGFR2 and NOTCH1 can undergo post-translational modifications, such as phosphorylation and ubiquitination, that affect their stability and activity. Crosstalk between these pathways can modulate these modifications, altering the signaling dynamics.

Implications for cancer therapy

Understanding the molecular interactions between FGFR2 and NOTCH1 provides new avenues for therapeutic intervention. Several strategies can be envisioned:

- Combination therapies:** Targeting both FGFR2 and NOTCH1 pathways simultaneously may produce synergistic effects, enhancing therapeutic efficacy and overcoming resistance to single-agent therapies.
- Biomarker development:** Identifying biomarkers that reflect the status of FGFR2 and NOTCH1 signaling can guide the selection of patients who are most likely to benefit from targeted therapies.
- Context-specific interventions:** Given the context-dependent roles of FGFR2 and NOTCH1 in cancer, therapies need to be tailored to the specific genetic and molecular landscape of each tumor.

Current therapeutic approaches

FGFR2 inhibitors: Several FGFR2 inhibitors are currently in clinical development or have received regulatory approval. These include small molecule tyrosine kinase inhibitors, monoclonal antibodies and FGF ligand traps. Clinical trials have shown promising results, particularly in cancers with FGFR2 alterations [5].

NOTCH1 inhibitors: NOTCH1 inhibitors, including γ -secretase inhibitors (GSIs) and monoclonal antibodies targeting NOTCH1 ligands, have been investigated in preclinical and clinical studies. While these inhibitors have shown efficacy in certain cancers, challenges such as toxicity and resistance need to be addressed.

*Address for Correspondence: Santiago Antonio, Department of Pathology, School of Medicine, National and Kapodistrian University of Athens, 11527 Athens, Greece; E-mail: antonio@santiago.uoa.gr

Copyright: © 2024 Antonio S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 20 April, 2024, Manuscript No. jmhmp-24-140977; Editor Assigned: 22 April, 2024, PreQC No. P-140977; Reviewed: 06 May, 2024, QC No. Q-140977; Revised: 13 May, 2024, Manuscript No. R-140977; Published: 20 May, 2024, DOI: 10.37421/2684-494X.2024.9.232

Challenges and future directions

Despite the promising therapeutic potential of targeting FGFR2 and NOTCH1, several challenges remain:

1. **Resistance mechanisms:** Cancer cells can develop resistance to FGFR2 and NOTCH1 inhibitors through various mechanisms, such as secondary mutations and activation of compensatory pathways. Understanding these resistance mechanisms is crucial for developing effective combination therapies.
2. **Toxicity:** Inhibition of FGFR2 and NOTCH1 signaling can lead to adverse effects, given their roles in normal tissue homeostasis. Strategies to minimize toxicity while maintaining therapeutic efficacy are needed.
3. **Biomarker identification:** Reliable biomarkers are essential for patient stratification and monitoring therapeutic responses. Advances in genomics and proteomics hold promise for identifying such biomarkers.

Conclusion

The intricate molecular interactions between FGFR2 and NOTCH1 have profound implications for cancer therapy. By elucidating the mechanisms of crosstalk and developing targeted interventions, we can improve therapeutic outcomes for patients with cancers driven by aberrant FGFR2 and NOTCH1 signaling. Future research should focus on overcoming the challenges of resistance, toxicity and biomarker identification to fully realize the potential of targeting these pathways in cancer therapy.

Acknowledgement

None.

Conflict of Interest

None.

References

1. Volpe, Virginia O., Daniel M. Klufas, Upendra Hegde and Jane M. Grant-Kels, et al. "The new paradigm of systemic therapies for metastatic melanoma." *J Am Acad Dermatol* 77 (2017): 356-368.
2. Ugurel, Selma, Joachim Röhmel, Paolo A. Ascierto and Keith T. Flaherty, et al. "Survival of patients with advanced metastatic melanoma: the impact of novel therapies—update 2017." *Eur J Cancer* 83 (2017): 247-257.
3. Tripathy, Nirmalya, Rafiq Ahmad, Seung Hyuck Bang and Gilson Khang, et al. "Outstanding antibiofilm features of quanta-CuO film on glass surface." *ACS Appl Mater Interfaces* 8 (2016): 15128-15137.
4. Piragine, Eugenia, Davide Petri, Alma Martelli and Vincenzo Calderone, et al. "Adherence to oral antidiabetic drugs in patients with type 2 diabetes: systematic review and meta-analysis." *J Clin Med* 12 (2023): 1981.
5. Kaufman, Howard L., Kim Margolin and Ryan Sullivan. "Management of metastatic melanoma in 2018." *JAMA Oncol* 4 (2018): 857-858.

How to cite this article: Antonio, Santiago. "Molecular Interactions between FGFR2 and NOTCH1: Implications for Cancer Therapy." *J Mol Hist Med Phys* 9 (2024): 232.