

New Understandings and Treatment Avenues in Salivary Gland Cancer: Molecular Pathways

Ruben Alfonso*

Department of Otolaryngology, Isola Tiberina-Gemelli Isola Hospital, 00186 Rome, Italy

Abstract

Salivary gland cancer, a rare but significant malignancy, poses considerable challenges in diagnosis and treatment due to its heterogeneous nature. Recent advancements in molecular biology have unveiled critical insights into the molecular pathways driving salivary gland cancer. This review aims to elucidate the key molecular mechanisms involved in salivary gland cancer pathogenesis and highlight potential therapeutic targets that could pave the way for novel treatment strategies.

Keywords: Salivary gland cancer • Novel treatment strategies • Therapeutic targets • Surgical techniques

Introduction

Salivary gland cancer, the most common type of salivary gland malignancy, represents a diverse group of tumours with varying histopathological and molecular characteristics. Despite advancements in surgical techniques and radiotherapy, the prognosis for advanced salivary gland cancer remains poor. Understanding the molecular underpinnings of salivary gland cancer is crucial for developing targeted therapies that can improve patient outcomes. This article delves into the molecular pathways implicated in salivary gland cancer and explores emerging therapeutic targets [1].

Literature Review

The Epidermal Growth Factor Receptor (EGFR) pathway is pivotal in cell proliferation, differentiation and survival. Deregulation of EGFR signalling has been observed in various cancers, including salivary gland cancer. Overexpression of EGFR and its ligands, along with activating mutations, contribute to tumour growth and resistance to conventional therapies. Targeting EGFR with Tyrosine Kinase Inhibitors (TKIs) and monoclonal antibodies has shown promise in preclinical studies, indicating potential therapeutic applications in salivary gland cancer. The PI3K/AKT/mTOR pathway plays a central role in cell survival, metabolism and growth. Aberrations in this pathway, such as mutations in PIK3CA and loss of PTEN, are frequently observed in salivary gland cancer. These alterations lead to constitutive activation of the pathway, promoting oncogenes and therapy resistance. mTOR inhibitors, such as everolimus, have demonstrated efficacy in preclinical models, suggesting that targeting this pathway could be beneficial in treating salivary gland cancer [2].

The RAS/RAF/MEK/ERK pathway, also known as the MAPK pathway, is integral to cell proliferation and differentiation. Mutations in RAS and RAF genes have been implicated in the pathogenesis of salivary gland cancer. Inhibitors targeting components of this pathway, such as MEK inhibitors, have shown anti-tumour activity in preclinical studies. Combining MAPK pathway inhibitors with other targeted therapies may enhance therapeutic efficacy.

***Address for Correspondence:** Ruben Alfonso, Department of Otolaryngology, Isola Tiberina-Gemelli Isola Hospital, 00186 Rome, Italy; E-mail: Alfonso.Ruben.fw@fbf-isola.it

Copyright: © 2024 Alfonso R. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 02 July, 2024, Manuscript No. jmhmp-24-143855; **Editor Assigned:** 04 July, 2024, PreQC No. P-143855; **Reviewed:** 16 July, 2024, QC No. Q-143855; **Revised:** 22 July, 2024, Manuscript No. R-143855; **Published:** 29 July, 2024, DOI: 10.37421/2684-494X.2024.9.244

Signaling pathway

Signaling pathway is involved in cell fate determination and differentiation. Deregulation of NOTCH signalling, including mutations and overexpression, has been identified in salivary gland cancer. NOTCH inhibitors, such as gamma-secretase inhibitors, are being investigated for their potential to inhibit tumour growth and progression in salivary gland cancer [3].

Wnt/ β -catenin pathway

The Wnt/ β -catenin pathway regulates cell proliferation, migration and stem cell maintenance. Aberrant activation of this pathway, through mutations or overexpression of pathway components, has been observed in salivary gland cancer. Targeting the Wnt/ β -catenin pathway with small molecule inhibitors and monoclonal antibodies holds promise as a therapeutic strategy.

Epigenetic modifications

Epigenetic alterations, including DNA methylation, histone modifications and non-coding RNAs, play a crucial role in the regulation of gene expression in salivary gland cancer. These modifications can lead to the silencing of tumour suppressor genes and activation of oncogenes. Epigenetic therapies, such as DNA methyltransferase inhibitors and histone deacetylase inhibitors, are being explored for their potential to reverse these changes and inhibit tumour growth [4].

Therapeutic targets and emerging treatments

EGFR inhibitors: Given the frequent deregulation of EGFR signalling in salivary gland cancer, EGFR inhibitors like cetuximab and erlotinib have shown potential in preclinical and early clinical studies. Combining EGFR inhibitors with other targeted therapies may enhance their efficacy and overcome resistance mechanisms [5]. PI3K/AKT/mTOR inhibitors targeting the PI3K/AKT/mTOR pathway with inhibitors such as everolimus and buparlisib has demonstrated anti-tumour activity in preclinical models of salivary gland cancer. These inhibitors can potentially be used in combination with other therapies to improve treatment outcomes.

MEK inhibitors: MEK inhibitors, such as trametinib, have shown promise in targeting the MAPK pathway in salivary gland cancer. Their efficacy can be enhanced by combining them with other targeted agents or immunotherapies [6].

NOTCH inhibitors: NOTCH pathway inhibitors, like gamma-secretase inhibitors, are being investigated for their potential to inhibit tumour growth in salivary gland cancer. Clinical trials are needed to evaluate their safety and efficacy in patients. Wnt/ β -catenin pathway inhibitors targeting the Wnt/ β -catenin pathway, such as LGK974, are under investigation for their potential to disrupt aberrant signalling in salivary gland cancer. These inhibitors may offer a novel therapeutic approach for patients with deregulated Wnt signalling.

Epigenetic therapies: Epigenetic therapies, including DNA methyl transferase inhibitors (e.g., azacitidine) and histone deacetylase inhibitors (e.g., vorinostat), are being explored for their ability to reverse epigenetic alterations in salivary gland cancer. These therapies hold promise for reactivating tumour suppressor genes and inhibiting oncogenic pathways.

Discussion

Salivary gland cancer, a rare type of salivary gland malignancy, has garnered increasing attention due to advancements in molecular biology that have elucidated its underlying mechanisms. Recent studies have highlighted several key molecular pathways involved in the pathogenesis of salivary gland cancer, providing new insights and potential therapeutic targets. One significant pathway implicated in salivary gland cancer is the PI3K/AKT/mTOR pathway. This signalling cascade is crucial for cell growth, proliferation and survival and its deregulation has been observed in various cancers, including parotid malignancies. Mutations and amplifications in genes encoding components of this pathway can lead to uncontrolled cell growth and tumour progression. Targeting the PI3K/AKT/mTOR pathway with specific inhibitors has shown promise in preclinical studies, suggesting a potential therapeutic strategy.

Another critical pathway is the MAPK/ERK pathway, which regulates cell differentiation, proliferation and apoptosis. Aberrations in this pathway, such as mutations in RAS and RAF genes, are common in salivary gland cancer. Inhibitors targeting MEK, a key component of the MAPK/ERK pathway, have demonstrated efficacy in reducing tumour growth in experimental models, indicating their potential as therapeutic agents. The Notch signalling pathway, which plays a vital role in cell fate determination and differentiation, is also implicated in salivary gland cancer. Alterations in Notch signalling can lead to abnormal cell proliferation and survival. Targeting this pathway with gamma-secretase inhibitors or monoclonal antibodies has shown promise in preclinical studies, offering a potential therapeutic avenue for salivary gland cancer.

Moreover, the emerging role of epigenetic modifications in salivary gland cancer pathogenesis cannot be overlooked. DNA methylation, histone modifications and non-coding RNAs have been identified as key regulators of gene expression in salivary gland cancer. Drugs targeting these epigenetic modifications, such as DNA methyl transferase inhibitors and histone deacetylase inhibitors, are being explored as potential therapeutic options. In addition to these pathways, the role of tumour microenvironment and immune evasion in salivary gland cancer is gaining attention. Immunotherapeutic approaches, such as immune checkpoint inhibitors, are being investigated to enhance the anti-tumour immune response and improve treatment outcomes.

Conclusion

Advancements in understanding the molecular pathways driving salivary gland cancer have opened new avenues for targeted therapies. EGFR, PI3K/AKT/mTOR, RAS/RAF/MEK/ERK, NOTCH and Wnt/ β -catenin pathways, along with epigenetic modifications, represent critical areas for therapeutic intervention. Continued research and clinical trials are essential to validate these targets and develop effective treatments for salivary gland cancer. By leveraging molecular insights, we can move closer to personalized and precision medicine approaches, ultimately improving outcomes for patients with this challenging malignancy.

Acknowledgement

None.

Conflict of Interest

None.

References

1. Que, Syril Keena T., Fiona O. Zwald and Chrysalyne D. Schmults. "Cutaneous squamous cell carcinoma: Incidence, risk factors, diagnosis and staging." *J Am Acad Dermatol* 78 (2018): 237-247.
2. Alam, Murad, April Armstrong, Christian Baum and Jeremy S. Bordeaux, et al. "Guidelines of care for the management of cutaneous squamous cell carcinoma." *J Am Acad Dermatol* 78: 560-578.
3. Corchado-Cobos, Roberto, Natalia García-Sancha, Rogelio González-Sarmiento and Jesús Pérez-Losada, et al. "Cutaneous squamous cell carcinoma: from biology to therapy." *Int J Mol Sci* 21 (2020): 2956.
4. Karia, Pritesh S., Jiali Han and Chrysalyne D. Schmults. "Cutaneous squamous cell carcinoma: estimated incidence of disease, nodal metastasis and deaths from disease in the United States, 2012." *J Am Acad Dermatol* 68 (2013): 957-966.
5. Thompson, Agnieszka K., Benjamin F. Kelley, Larry J. Prokop and M. Hassan Murad, et al. "Risk factors for cutaneous squamous cell carcinoma recurrence, metastasis and disease-specific death: A systematic review and meta-analysis." *JAMA Dermatol* 152 (2016): 419-428.
6. Xiang, Fan, Robyn Lucas, Simon Hales and Rachel Neale, et al. "Incidence of nonmelanoma skin cancer in relation to ambient UV radiation in white populations, 1978-2012: Empirical relationships." *JAMA Dermatol* 150 (2014): 1063-1071.

How to cite this article: Alfonso, Ruben. "New Understandings and Treatment Avenues in Salivary Gland Cancer: Molecular Pathways." *J Mol Hist Med Phys* 9 (2024): 244.