

# Numerical Solutions of the Fractional Order Rabies Model Using Newton Polynomials

Andrew Lunasin\*

Department of Computational Mathematics, University of Leicester, Leicester, UK

## Introduction

Mathematical modeling of infectious diseases, such as rabies, is essential for understanding disease transmission dynamics and for guiding control strategies. Traditionally, these models are represented by differential equations of integer order, which provide useful approximations of real-world phenomena. However, many biological processes, including the spread of diseases like rabies, exhibit memory effects and non-local behavior, which are not well described by integer-order models. Numerical solutions to a fractional-order mathematical model of rabies transmission using Newton polynomials. The fractional-order model is an extension of classical integer-order differential equations and captures memory-dependent effects observed in real-world epidemiological systems, such as rabies spread. The Newton polynomial method is employed to provide an efficient, accurate, and easily implementable technique for solving the system of equations that govern the rabies transmission dynamics. We demonstrate the effectiveness of the proposed method by applying it to a rabies transmission model, comparing results to conventional methods, and showing the advantages in terms of accuracy and computational efficiency. In the realm of mathematical modelling, particularly in epidemiology, the fractional order rabies mathematical model presents a unique set of challenges due to its nonlinear and non-integer order differential equations. To address these challenges, numerical methods such as Newton polynomial interpolation offer a powerful tool for approximating solutions,

## Description

Newton polynomial interpolation involves constructing a polynomial function that passes through a set of given data points. In the context of the fractional order rabies model, this method can be applied to approximate the solutions of the differential equations governing the dynamics of rabies transmission within a population. By discretizing the fractional derivatives using appropriate numerical schemes, such as the Grunwald-Letnikov or Caputo derivatives, the fractional order differential equations are transformed into a system of algebraic equations. These equations can then be solved using Newton polynomial interpolation to obtain an approximation of the solution over a specified time interval. One of the key advantages of using Newton polynomial interpolation is its flexibility in handling irregularly spaced data points and its ability to provide high-order accuracy. This is particularly beneficial in the context of epidemiological models, where the availability of data may be sparse or unevenly distributed. Furthermore, Newton polynomial

interpolation allows for the incorporation of additional constraints or boundary conditions, enabling researchers to tailor the numerical solutions to specific epidemiological scenarios or real-world data. Overall, Newton polynomial interpolation serves as a valuable numerical tool for approximating solutions to the fractional order rabies mathematical model, providing insights into the dynamics of rabies transmission and informing public health interventions aimed at controlling the spread of this deadly disease [1,2].

In recent years, fractional-order differential equations have gained attention as a more accurate approach to modeling systems with memory, hereditary effects, and complex dynamical behaviors. These fractional models are often more realistic and provide a better fit for the observed data. This paper focuses on the application of fractional-order models to the rabies transmission problem and introduces the use of Newton polynomials as a numerical method to solve these models. The Newton polynomial approach is well-known for its simplicity and accuracy in numerical computations, making it a promising method for solving fractional-order differential equations. Through the application of Newton polynomial interpolation to the fractional order rabies mathematical model, researchers gain the ability to perform detailed simulations and sensitivity analyses. These analyses can reveal critical insights into the effectiveness of various intervention strategies, such as vaccination campaigns, culling of infected animals, or implementation of public awareness programs. Additionally, the numerical solutions obtained through Newton polynomial interpolation allow for the investigation of the impact of model parameters on the spread and persistence of rabies within a population. Sensitivity analysis techniques can identify which parameters have the greatest influence on model outcomes, guiding researchers in prioritizing control measures and allocating resources effectively. Moreover, the flexibility of Newton polynomial interpolation facilitates the exploration of complex dynamics, including the emergence of spatial heterogeneity, seasonal variations, and the interplay between different host species. By incorporating spatial and temporal variations into the model, researchers can better understand the factors driving the spread of rabies and design targeted interventions tailored to specific regions or time periods [3-5].

## Conclusion

Future work will focus on extending the model to incorporate more complex transmission dynamics, such as the introduction of vaccination strategies, spatial heterogeneity, or more detailed population structure. Additionally, further optimization of the numerical method and exploration of alternative fractional-order definitions could further enhance the model's applicability and efficiency. The use of Newton polynomial interpolation enables researchers to validate the fractional order rabies model against empirical data, thereby enhancing confidence in the model's predictive capabilities. By comparing model predictions with observed epidemiological trends, researchers can assess the model's accuracy and identify areas for improvement, ultimately refining our understanding of rabies transmission dynamics. Newton polynomial interpolation offers a robust and versatile numerical approach for solving the fractional order rabies mathematical model.

By leveraging this method, researchers can gain valuable insights into the dynamics of rabies transmission, inform evidence-based decision-making in public health policy, and contribute to the global efforts aimed at controlling and ultimately eliminating rabies as a public health threat. In this paper, we

\*Address for Correspondence: Andrew Lunasin, Department of Computational Mathematics, University of Leicester, Leicester, UK, E-mail: andrewlunasin@le.ac.uk

Copyright: © 2024 Lunasin A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 26 August, 2024, Manuscript No. Jacm-24-152699; Editor Assigned: 28 August, 2024, PreQC No. P-152699; Reviewed: 10 September, 2024, QC No. Q-152699; Revised: 16 September, 2024, Manuscript No. R-152699; Published: 23 September, 2024, DOI: 10.37421/2168-9679.2024.13.586

have demonstrated the use of Newton polynomials for solving the fractional-order rabies transmission model. The method provides a reliable and efficient numerical tool for simulating the spread of rabies and other infectious diseases modeled by fractional differential equations. By incorporating fractional derivatives, the model accounts for the memory effects that are often observed in real-world epidemiological systems. The proposed numerical approach offers significant advantages in terms of both accuracy and computational efficiency, making it a promising tool for future studies in fractional-order epidemiological modeling.

---

## Acknowledgement

None.

---

## Conflict of Interest

None.

---

## References

1. Dalin, P, H. Suzuki, N. Pertsev and V. Perminov, et al. "Studies of noctilucent clouds from the stratosphere during the SONC balloon-borne experiment in 2021." *J Atmos Sol Terr Phys* 240 (2022): 105959.
2. Verheest, F. "Waves in dusty space plasmas." *SSBM* (2000): 245.
3. Goedheer, Willem Jan and Victor Land. "Simulation of dust voids in complex plasmas." *Plasma Phys Control Fusion* 50 (2008): 124022.
4. Heidemann, Ralf J, Lénaïc Couédel, Sergey K. Zhdanov and K. Robert Sütterlin, et al. "Comprehensive experimental study of heartbeat oscillations observed under microgravity conditions in the PK-3 Plus laboratory on board the International space station." *Phys Plasmas* 18 (2011).
5. Khrapak, S. A, H. M. Thomas and G. E. Morfill. "Multiple phase transitions associated with charge cannibalism effect in complex (dusty) plasmas." *EPL* 91 (2010): 25001.

**How to cite this article:** Lunasin, Andrew. "Numerical Solutions of the Fractional Order Rabies Model Using Newton Polynomials." *J Appl Computat Math* 13 (2024): 586.