

The Progression of a Treatable Illness

Billey Maccum*

Institute for Biomedical Ethics and History of Medicine, University of Zurich, Zurich, Switzerland

Description

A vulnerable energy-consuming renal tubule and a hypoxic, hyperosmolar microenvironment emerged from evolutionary adaptation to a terrestrial setting. Natural selection supports a successful energy investment strategy: during reproductive years, energy is committed to maintaining nephron integrity, but this decreases with increasing senescence after w40 years of age. Restricted foetal growth or preterm delivery (a life history trade-off that results in fewer nephrons), evolutionary selection for APOL1 mutations (a trade-off that provides resistance to trypanosome infection), and modern life experience are all risk factors for chronic kidney disease (Western diet mismatch leading to diabetes and hypertension). Current advancements in genomes, epigenetics, and developmental biology have identified proximal reasons of kidney disease, but slowing renal disease remains a challenge. By addressing the root causes of kidney illness, evolutionary medicine offers a supplementary solution. Evolutionary mechanisms are responsible for significant variations in nephron number at birth, nephron heterogeneity, and changing vulnerability to kidney injury throughout time [1]. New techniques could emerge from combining molecular genetics, evolutionary developmental biology (evo-devo), developmental programming, and life history theory.

Nephron loss, hyperfiltration, and incomplete repair are all symptoms of progressive kidney disease, which is referred to as a "maladaptive" process. Evolutionary medicine is a new science that has evolved in the last 20 years and has broadened study frontiers. Evolutionary adaptation, unlike physiologic (homeostatic) adaptation, is the result of reproductive success and natural selection. Mismatch of the phenotype with the environment or evolutionary trade-offs can both lead to evolutionary explanations for physically maladaptive responses [2].

The global epidemic of Chronic Kidney Disease (CKD) has motivated new techniques to unravel the underlying mechanisms: the incidence of End-stage Kidney Disease (ESKD) in children is 10 per million, whereas the lifetime risk of ESKD in adults is more than 5%. Diabetic nephropathy is now the most common cause of ESKD in adults, whereas congenital abnormalities of the kidneys and urinary tract (CAKUT) are the most common cause in children. More recently, epidemiologic studies have revealed that incomplete recovery from acute kidney damage (AKI) is a risk factor for developing CKD, and that CKD enhances susceptibility to AKI: the proximal tubule thus becomes a key target of injury and CKD progression. The gradual understanding of kidney anatomy and function, as well as the fundamental role of the nephron in maintaining homeostasis, has led to advances in nephrology [3,4]. Jean Oliver's elegant morphologic studies in the early twentieth century revealed the development of intermixed hypertrophied and atrophied nephrons in the

kidneys of CKD patients, which was a significant finding. Oliver also described the widespread production of atubular glomeruli and aglomerular tubules in the kidneys of patients with severe CKD using microdissection techniques. The emergence of atubular glomeruli as a result of proximal tubular damage is a hallmark of late CKD, according to morphometric investigations of kidneys from patients with CKD due to vascular, glomerular, tubulointerstitial, or toxic etiologies [5].

Systems biology could provide fresh evolutionary insights into CKD maladaptation by merging physiologic and genomic approaches across vast sample sizes of numerous species' life histories. New medicines could be developed to target the processes that cause the population of cells in diseased or wounded tissue to be selected. The clues to renal senescence may be discovered in the epigenetic mechanisms of foetal programming, accounting for the transition from paediatric to adult CKD, if kidney research is framed in the perspective of life history theory.

Acknowledgements

None.

Conflict of Interest

There are no conflicts of interest by author.

References

1. Shimizu, Masaki, Keita Katayama, Eiji Kato and Shiro Miyayama, et al. "Evolution of acute focal bacterial nephritis into a renal abscess." *Pediatr Nephrol* 20 (2005): 93-95.
2. Abudayyeh, Ala A., Amit Lahoti and Abdulla K. Salahudeen. "Onconephrology: The need and the emergence of a subspecialty in nephrology." *Kidney Int* 85 (2014): 1002-1004.
3. Ghiggeri, Gian M., Monica Dagnino, Stefano Parodi and Cristina Zennaro, et al. "Discordant evolution of nephrotic syndrome in mono- and dizygotic twins." *Pediatr Nephrol* 21 (2006): 419-422.
4. Hecht, Natalie, Abiodun Omolaja, Dave Witte and Leonardo Canessa. "Evolution of antiglomerular basement membrane glomerulonephritis into membranous glomerulonephritis." *Pediatr Nephrol* 23 (2008): 477-480.
5. Wang, Angela Y.M., Yu An, Guang-Yan Cai and Jiang-Hua Chen, et al. "Nephrology in China." *Nephrol Worldwide* (2021): 251-290.

*Address for Correspondence: Billey Maccum, Institute for Biomedical Ethics and History of Medicine, University of Zurich, Zurich, Switzerland; E-mail: mac.billey@yahoo.com

Copyright: © 2022 Maccum B. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 02 April, 2022, Manuscript No. JNT-22-62524; Editor Assigned: 04 April, 2022, PreQC No. P-62524; Reviewed: 16 April, 2022, QC No. Q-62524; Revised: 21 April, 2022, Manuscript No. R-62524; Published: 28 April, 2022, DOI: 10.37421/2161-0959.2022.12.387

How to cite this article: Maccum, Bille. "The Progression of a Treatable Illness." *J Nephrol Ther* 12 (2022): 387.